

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
**09.04.2003 Bulletin 2003/15**

(21) Application number: **02079804.7**

(22) Date of filing: **21.04.1992**

(51) Int Cl.<sup>7</sup>: **A23K 1/17, A23L 3/3463,**  
**A23L 3/3571, C12N 1/06,**  
**C12N 1/15, C12N 1/19,**  
**C12N 1/20, C12N 7/01,**  
**C12N 9/14, C12N 15/33,**  
**C12N 15/55, C12N 15/74,**  
**C12Q 1/04, C12Q 1/10,**  
**C12Q 1/18, C12Q 1/34,**  
**C12Q 1/70**  
**// C12R1:145**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GR IT LI LU MC NL PT SE**

(30) Priority: **20.04.1991 GB 9108498**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**92303533.1 / 0 510 907**

(71) Applicant: **AGRICULTURAL AND FOOD**  
**RESEARCH COUNCIL**  
**Colney, Norwich NR4 7UA (GB)**

(72) Inventor: **Gasson, Michael John**  
**Dereham, Norfolk NR20 5NA (GB)**

(74) Representative: **Thomas, Philip John Duval et al**  
**Eric Potter Clarkson,**  
**Park View House,**  
**58 The Ropewalk**  
**Nottingham NG1 5DD (GB)**

Remarks:

This application was filed on 18 - 11 - 2002 as a  
divisional application to the application mentioned  
under INID code 62.

(54) **Lysin of a bacteriophage of Clostridia spp**

(57) Bacteriophages of food-contaminating or patho-  
genic bacteria or the lysine thereof are used to kill such  
bacteria. Examples include lysins from bacteriophages  
of *Listeria monocytogenes* and *Clostridium tyrobutyri-*  
*cum*.

Tests for bacterial contamination can be made spe-  
cific for specific bacteria by using the appropriate bac-  
teriophage or lysin thereof and determining whether  
cells are lysed thereby.



Fig. 1

## Description

[0001] This invention relates to the use of bacterial viruses (bacteriophages) which use bacteria as hosts and produce a bacteriophage lysin responsible for cell-wall degradation and lysis of the host cells.

[0002] Attempts to use a bacteriophage as an antimicrobial agent have failed to be effective. We have previously used the lysin of the bacteriophage  $\phi$ vML3 of *Lactococcus lactis* ML3, which is active against all strains of all subspecies of *Lactococcus lactis*, very weakly affects group D enterococci, but does not have any action on a wide variety of other species tested (Shearman et al (1989) *Molecular and General Genetics* 218: 214-221), to lyse cheese starter cultures (WO90/00599). WO90/00599 also discloses the use of micro-organisms, transformed to express the  $\phi$ vML3 lysin, to suppress populations of bacteria susceptible to the lysin, ie the *Lactococcus lactis* cheese starter culture strains.

[0003] It is also known to use cheese starter culture bacteria to produce the simple peptide nisin in order to destroy harmful bacteria.

[0004] We have now found that further bacteriophage lysins can be used to destroy unwanted bacteria, especially food-contaminating bacteria prejudicial to health.

[0005] A first aspect of the present invention provides a formulation comprising a lysin of bacteriophage  $\phi$ LM4 or Clostridia bacteriophage, or a variant thereof having an amino acid sequence which is substantially identical to such a lysin, said formulation being substantially free of the bacteriophage itself.

[0006] Preferably the Listeria phage  $\phi$ LM4 or Clostridium tyrobutyricum phage  $\phi$ P1 lysins are used. They act against all tested species and strains of Listeria and also strains of Kurthia zopfii, or against Clostridium tyrobutyricum (as appropriate), but lack activity against other tested species.

[0007] A "variant" of such a lysin is any polypeptide of which at least 30% (preferably at least 50%, 75%, 90%, 95% or 99%) has at least 80% (preferably at least 90%, 95% or 99%) amino acid homology with the corresponding region of the lysin itself and which has at least 30% (preferably at least 50%, 75%, 90% or 95%) of the bacterial lysing capability of the said lysin.

[0008] Food-contaminating bacteria are those which, by virtue of their presence or compounds produced by them, cause undesirable flavours, odours or visual appearances or cause illness in humans or animals consuming the food.

[0009] The organism which is destroyed may be any of the following:

[0010] Listeria monocytogenes, Clostridium tyrobutyricum, Clostridium botulinum, Clostridium perfringens, lactic acid bacteria (eg Lactobacillus brevis) causing beer spoilage, Salmonella spp., Yersinia spp., Campylobacter, E. coli, Pseudomonas spp., Staphylococcus, Bacillus spp. (including Bacillus cereus), Shigella spp. and Vibrio spp.

[0011] Pathogenic bacteria include all pathogenic bacteria of humans, animals and plants. However, in a medical or veterinary context, as is explained further below, bacteria involved in topical or superficial infections are of particular interest. These include Staphylococcus spp. (eg Staph. aureus), Streptococcus spp., Corynebacterium spp., Clostridium spp. (eg Cl. perfringens), Yersinia spp. (eg Y. pestis), Pasteurella spp. (eg P. multocida), Streptobacillus spp. (eg Streptobacillus moniliformis), Proteus spp. (eg P. mirabilis) and Pseudomonas spp.

[0012] A second aspect of the invention provides a substantially pure preparation of a lysin of bacteriophage  $\phi$ LM4 or a variant thereof having an amino acid sequence which is substantially identical to such a lysin.

[0013] A third aspect of the invention provides a nucleotide coding sequence for a lysin included in the formulation of the invention.

[0014] A fourth aspect provides an expression vehicle comprising a coding sequence according to the invention and regulatory regions associated therewith for expression of the coding sequence in a suitable host.

[0015] Suitable regulatory expression vectors, transformation techniques, and hosts are all known in the art. The host may be any micro-organism or cell line which is found to express the said lysin gene, and may be a bacterium such as E. coli or Lactococcus lactis, a yeast such as Saccharomyces cerevisiae or Kluyveromyces fragilis or a filamentous fungus such as Aspergillus niger.

[0016] Thus a fifth aspect provides a microbial or cell line host transformed with means to express a lysin, said means comprising the coding sequence of the invention.

[0017] A sixth aspect provides a lysin derived from the cultivation of such a host.

[0018] A seventh aspect of the present invention provides a method of destroying micro-organisms characterized in that said micro-organisms are lysed with a formulation of the invention.

[0019] The use of the invention in food or agriculture simply involves the addition of an amount sufficient to provide an inhibitory concentration of lysin activity. The specific activity of any preparation may readily be calculated, for example by use of the spectrophotometric assay described later. The quantity of preparation necessary for effective protection in a given food may be arrived at by routine experimentation. The lysin is applied in a suitable, nontoxic aqueous medium. Any food may be treated with such a preparation by addition or application to surfaces eg of cut, cooked meat or poultry, soft cheeses and pâtés of fish or meat. The term "food" includes drinks (such as water, beer, milk and soft drinks), animal food (such as pet food or cattle food) and produce destined for consumption by humans or animals (such as stored potatoes). In agriculture, a particular application is addition to silage where Listeria and Clostridium

tyrobutyricum are known to present a problem that can be passed on up the food chain. In brewing, brewing yeast transformed with a lysin gene may be used.

[0020] In a medical or veterinary context, because the lysin is likely to be degraded or to produce an immune reaction, it is preferred to administer it topically in diseases of the skin such as ulcers, burns and acne. It may be applied as the clinician directs, as a lotion, cream or ointment.

[0021] An eighth aspect provides a method of testing for Clostridia in a sample, comprising adding bacteriophage lysin substantially free of intact bacteriophage to the sample and determining whether bacterial cells have been lysed thereby, wherein the lysin is a lysin of Clostridial bacteriophage or a variant thereof having an amino acid sequence which is substantially identical to such a lysin.

[0022] Any technology that exploits the release of intracellular biochemicals (eg ATP or enzymes such as alkaline phosphatase or esterase) to detect micro-organisms can, in accordance with the invention, be made specific for the target range of such lysins. For example, an ATP or phosphatase release test for *Listeria* using the *Listeria* bacteriophage or lysin thereof, in which the release of ATP or phosphatase is detected (eg by linkage to a luciferase reaction and monitoring of photon release or by spectrophotometric methods as is described below) indicates the specific presence of *Listeria* in a sample. The invention further provides a kit comprising a lysin and means to detect bacterial lysis.

[0023] Preferably, the bacteriophage in all these contexts is or at least includes *Listeria monocytogenes*  $\phi$ LM4 or a bacteriophage of *Clostridium tyrobutyricum*, such as  $\phi$ P1. Several different lysins may be used in order to destroy or identify a specific range of bacteria.

[0024] The cloning and characterization of the gene for the lysin of the *Listeria* bacteriophage  $\phi$ LM4 has facilitated the production of the free lysin and the availability of its structural gene. These components have application in the protection of environment and food material from pathogenic strains of *Listeria*. The free lysin acts as a novel antimicrobial that kills such bacteria and the gene can be genetically engineered in a non-pathogenic micro-organism such that the latter produce the *Listeria* lysin thereby equipping it with a novel anti-*Listeria* capability. For example, a food-grade micro-organism may be transformed with a DNA construct comprising a coding sequence for the lysin.

[0025] Preferred embodiments of the invention will now be described by way of example with reference to the accompanying drawings, in which:-

Figure 1 shows patches of *E. coli* clones with *Hind*III fragments of  $\phi$ LM4 DNA in the *Hind*III site of vector pUC18. The plate is overlaid with a suspension of *Listeria monocytogenes* 6868 cells and lysin producing clones create clear zones around the patch (indicated by an arrow).

Figure 2 is a restriction and deletion map of lysin-expressing clone pF1322. The result of lysin activity tests is indicated to the right. The inferred location of the lysin gene is shown. Arrows indicate the orientation of the lysin gene with respect to the *lac*  $\alpha$  promoter of the pUC vector used which is transcribed from left to right in this figure (ie pF1324 is opposed to the *lac*  $\alpha$  promoter, other clones are transcribed in the same direction as the *lac*  $\alpha$  promoter).

Plasmid pF1322 is pUC18 carrying a 3.6kb *Hind*III fragment of bacteriophage  $\phi$ LM4 DNA. Plasmid pF1326 is pF1322 with a 0.56kb *Hind*III - *Sal*I deletion. Plasmid pF1327 is pF1322

with a 1.32kb *Hind*III - *Eco*RI deletion. Plasmid pF1324 is pUC18 carrying a 1.9kb *Hind*III - *Nru*I fragment of pF1322 cloned between its *Hind*III and *Hinc*II sites. Plasmid pF1325 is pUC18 carrying a 1.6kb *Nru*I - *Hind*III fragment of pF1322 cloned between its *Hinc*II and *Hind*III sites. Plasmid pF1328 is pUC19 carrying a 1.9kb *Hind*III - *Nru*I fragment of pF1322 cloned between its *Hind*III and *Hinc*II sites. Plasmid pF1329 is pF1328 carrying a 1.6kb *Ba*PI deletion from the polylinker *Bam*HI site. Plasmid pF1330 is pF1328 carrying a 1.6kb *Ba*PI deletion from the polylinker *Bam*HI site.

Figure 3 illustrates the response of a suspension of *Listeria monocytogenes* 6868 cells to cell free extracts of *E. coli* strains harbouring plasmids pF1322( $\blacktriangle$ ), pF1328( $\Delta$ ), pF1329(o) and pUC19( $\circ$ ).

Figure 4 is a Coomassie blue stained SDS polyacrylamide gel of proteins produced by *E. coli* strain carrying the T7 expression vector pSP73 (tracks 2 and 3) or pF1331 which carries the lysin gene (tracks 4 and 5). Uninduced cells (tracks 2 and 4) are compared with induced cells (tracks 3 and 5). Molecular weight markers are present (tracks 1 and 6) and the expressed lysin protein is indicated by an arrow.

Figure 5 illustrates the sequencing strategy used. The extent and direction of sequences determined are indicated by the arrows. Synthetic oligonucleotide primers are indicated by boxes.

Figure 6 shows a single strand of the region of  $\phi$ LM4 DNA that encodes the lysin gene.

Figure 7 is the Analyseq print out of the analysis of the DNA sequence shown in Figure 6. The identification of the open reading frame of the lysin gene is in the top panel.

Figure 8 shows the double stranded DNA sequence of the lysin structural gene and its translated protein product.

Figure 9 shows the protective effect of cloned *Listeria* lysin on skimmed milk to which *Listeria Monocytogenes* is added.

Figure 10 shows the expression of the *Listeria* lysin gene in *Lactococcus lactis* under the control of the lactose inducible lactococcal lactose operon promoter.

#### EXAMPLE 1: CLONING OR LYSIN GENE, ETC

##### Isolation of bacteriophage $\phi$ LM4

[0026] A bacteriophage named  $\phi$ LM4 was isolated from a culture of *Listeria monocytogenes* serotype 4b that was originally obtained from a listeriosis outbreak in Nova Scotia, Canada in 1981. The source of the infection was tracked down to contaminated coleslaw. This culture of *Listeria monocytogenes* was deposited under the Budapest Treaty as NCTC 12452 in the National Collection of Type Cultures, Central Public Health Laboratory, Colindale, London, UK on 21 March 1991. The bacteriophage was purified by standard single plaque isolation procedure using *Listeria monocytogenes* F6868 as the host. This culture was similarly deposited under the Budapest Treaty as NCTC 12453 in the National Collection of Type Cultures, Central Public Health Laboratory, Colindale, London, UK on 21 March 1991. Examination of this bacteriophage by electron microscopy revealed it to have an isometric head with a diameter of approximately 50nm and a tail of approximately 250nm.

##### Isolation of DNA from *Listeria monocytogenes* bacteriophage $\phi$ LM4

[0027] 20ml of an 18 hour culture of *Listeria monocytogenes* F6868 was inoculated into 500ml of Bacto tryptose phosphate broth and incubated with shaking at 30°C. When O.D. 600 reached 0.15 the culture was infected with  $5 \times 10^{10}$  p.f.u. of bacteriophage  $\phi$ LM4 and incubated until lysis was apparent as a loss of turbidity. The lysate was centrifuged at 6000 x g for 10 min at 4°C. The bacteriophage lysate was then concentrated by polyethylene glycol precipitation and purified on caesium chloride stepped gradients using well established protocols (Bachrach and Friedmann (1971) *Applied Microbiology* 22: 706-715). Bacteriophage DNA was extracted by dialysis against 50% formamide in TE buffer (0.1 Tris - HCl, 0.01M EDTA, pH 8.5). Further purification was then performed on caesium chloride-ethidium bromide equilibrium density gradients. Examination of the bacteriophage DNA by agarose gel electrophoresis revealed the genome to be approximately 39kb in size.

##### Cloning the bacteriophage $\phi$ LM4 lysin gene

[0028] DNA purified from bacteriophage  $\phi$ LM4 was digested with restriction endonuclease *Hind*III and ligated to plasmid pUC18 vector DNA that had also been cleaved with restriction endonuclease *Hind*III. The ligated DNA was transformed into *Escherichia coli* TB1 and ampicillin resistant colonies were selected on LB agar containing 50µg/ml ampicillin, 40µg/ml isopropyl-β-D-thiogalactopyranoside (IPTG) and 40µg/ml 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-gal). These steps were performed using well established protocols (Sambrook, J. et al (1989), *Molecular Cloning. A laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 2nd Edition).

[0029] White colonies were screened for their ability to produce a bacteriophage lysin active against *Listeria monocytogenes*. These colonies were patched onto duplicate Bacto tryptose agar plates and incubated for 18 hours at 37°C. One plate was exposed to chloroform vapour for 10 min and then seeded with 0.2ml of an 18 hour broth culture of *Listeria monocytogenes* F6868. After incubation at 30°C for 18 hours clear zones of lysins were apparent around patches of clones expressing the *Listeria* bacteriophage  $\phi$ LM4 lysin. This is illustrated in Figure 1. Positive clones were recovered from the duplicate plate and the pUC18 derivative plasmid isolated and characterized by digestion with restriction endonuclease *Hind*III. One lysin expressing pUC18 clone that contained a 3.6kb insert of  $\phi$ LM4 DNA was chosen for further analysis. This plasmid was designated pF1322.

##### Deletion analysis of lysin expressing plasmid pF1322

[0030] Characterization of pF1322 was undertaken by constructing a restriction map of this insert using single and double digests with a variety of restriction enzymes. The map is presented in Figure 2. Deletion of some regions of the

3.6kb insert contained in pFI322 was achieved by digestion with certain of these enzymes, religation and transformation into *E. coli* TB1. In other instances endonuclease *Bal* 31 was used to introduce deletions. In addition, some regions of the 3.6kb cloned DNA in pFI322 were deleted by digestion with certain restriction endonucleases and re-cloning into appropriately cleaved plasmid vectors pUC18 or pUC19 and transformation into *E. coli* TB1. These manipulations are clearly documented in Figure 2 which is presented in the form of a deletion map for pFI322. After confirming that the various constructed plasmids derived from pFI322 had the expected structures, these clones were tested for their ability to produce *Listeria* bacteriophage lysin. As well as the plate assay described above and illustrated in Figure 1, a spectrophotometric assay was also used. For this the *E. coli* strain carrying plasmid clones were grown at 37°C for 18 hours, harvested by centrifugation at 6000 x g for 5 min at 4°C, washed down once in 100mM Tris buffer pH7.5 and resuspended in this same buffer at approximately 10mg dry weight/ml. Cell free extracts were made by 6 cycles of ultrasonication (15 sec on, 10 sec off) at 0°C using the microprobe of an MSE Soniprep™ 150. Unbroken cells and cell debris were removed by centrifugation at 25000 x g for 15 min at 4°C.

[0031] Samples of the cell free extracts were added to an equilibrated (5 min at 37°C) 4ml reaction mixture containing 400µmole Tris HCl pH7.5 and *Listeria monocytogenes* F6868 indicator cells that had been harvested and resuspended at an O.D. 600 of 2.3. The fall in optical density caused by lysis of indicator cells was followed using a spectrophotometer. Typical results from use of this protocol are presented in Figure 3. The lytic activity of the plasmid derivative described above and in Figure 2 were assessed using both of these methods and the results are presented in Figure 2.

[0032] These results demonstrated that the structural gene for bacteriophage φLM4 was contained within the left hand 1.2kb of the DNA cloned in pFI322 and defined by the *Hind*III site at co-ordinate 0 and the *Eco*R1 site at co-ordinate 1.25 of the map illustrated in Figure 2.

[0033] Figure 2 also indicates the orientation of *Listeria* bacteriophage φLM4 DNA with respect to the *E. coli* *lac* α promoter that is present on vectors pUC18 and pUC19. It is apparent that a positive reaction in the lysin assay is only found when one orientation is maintained (eg pFI324 is negative whereas pFI328 is positive even though both constructs contain the same *Listeria* bacteriophage φLM4 fragment). This suggests that expression of the lysin gene depends on use of the *E. coli* *lac* α promoter and that no *Listeria* bacteriophage φLM4 promoter is present and active in *E. coli*.

#### Detection of the lysin protein

[0034] In order to identify a protein produced by the fragment of φLM4 DNA that expressed lysin activity another *E. coli* vector was used. A 2kb fragment from plasmid pFI328 between the *Hind*III site at co-ordinate 0 and a unique *Bam*HI site present on the polylinker of pUC19 was isolated and cloned between the *Hind*III and *Bam*HI sites of the T7 expression vector pSP73 that was purchased from Promega. The constructed plasmid named pFI331 was transformed into the *E. coli* host strain JM109DE3.

[0035] The *E. coli* T7 promoter in this vector is expressed by the phage specific T7 RNA polymerase which is induced by addition of IPTG in the appropriate host strain *E. coli* JM109 DE3. Cultures of this strain carrying pSP73 as a control or pFI331 were grown for 3 hours and induced by addition of IPTG to a final concentration of 0.2mM. Incubation was continued for a further 3 hours before the cultures were harvested and used to prepare cell extracts using well-established, published procedures (Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorff, J.W. (1990) *Methods in Enzymology* 185: 60-89).

[0036] Proteins present in cell extracts were analysed using conventional SDS-polyacrylamide gel electrophoresis (Laemmli (1970) *Nature* 227: 680-685). The results presented in Figure 4 clearly demonstrate that the 2kb fragment of pFI331 expresses a single protein with a molecular size of 31 kilodaltons which represents the lysin enzyme.

#### DNA sequence of the *Listeria* bacteriophage φLM4 lysin gene

[0037] The region of DNA between co-ordinate 0 and 1.2 in Figure 2 was subject to oligonucleotide sequence analysis using the dideoxy chain-termination method (Sanger, F., Coulson, A.R., Barrell, B.G., Smith, A.J.H. and Roe, B.A. (1980) *J. Molec. Biol.* 143) with a sequenase™ version 2.0 kit (United States Biochemical Corporation). The 0.9kb *Hind*III - *Eco*RI and the 0.3kb *Eco*RI - *Eco*RI fragments of pFI328 were subcloned in the M13 sequencing vectors M13mp18 and M13mp19 to create templates and sequenced using universal and synthetic oligonucleotide primers. To sequence across the internal *Eco*RI site at co-ordinate 0.9 double stranded sequencing of pFI329 plasmid DNA was used. The sequencing strategy is presented in Figure 5 and the complete DNA sequence is in Figure 6. The sequence was analysed using the computer programme ANALYSEQ™ (Staden (1980) *Nucleic Acid Research* 8: 3673-3694) which revealed an open reading frame that represents the *Listeria* bacteriophage lysin gene. The printout from the Analyseq analysis is presented in Figure 7 and the open reading frame representing the lysin structural gene and its translated protein product is presented in Figure 8. The molecular size of the translated protein was calculated to be 32.9 kilodaltons which agrees well with the calculated 31 kilodalton size of the protein expressed by the T7 vector

pSP73 (Clone pFI331 in Figure 4).

Activity and specificity of the *Listeria* bacteriophage  $\phi$ LM4 lysin

[0038] Figure 3 illustrates the lytic activity of crude cell free extracts of *E. coli* TB1 carrying the plasmids pFI322, pFI328, pFI329 and pUC19 assayed using the spectrophotometric method described above. This activity was related to units of commercially available mutanolysin (Sigma) as has been described previously (Shearman, C., Underwood, H, Jury, K. and Gasson M. (1989) *Mol. Gen. Genetics* 218: 214-221). The crude cell extracts of lysin expressing clones typically contained 5000 mutanolysin equivalent units per mg. protein.

[0039] In order to test the spectrum of activity of this lysin, the spectrophotometric assay was performed on 16 serotypes of *Listeria monocytogenes*, all other species of *Listeria*, the related species *Kurthia zopfii* and a variety of other gram positive and gram negative bacteria. The results compiled in Table 1 show that the *Listeria* bacteriophage  $\phi$ LM4 lysin was active against all tested strains of *Listeria monocytogenes*, *Listeria innocua*, *Listeria ivanovii*, *Listeria murrayi*, *Listeria seelegri*, *Listeria welshimeri*, *Listeria grayi* and *Kurthia zopfii*. No activity was found against any of the other species tested.

TABLE 1:

ACTIVITY OF CLONED LYSIN AGAINST LISTERIA SPECIES				
Organism	Strain	Serotype	Relative <sup>a</sup> Activity	Time (min) <sup>b</sup> $\Delta OD_{600}=1$
<i>Listeria monocytogenes</i>	F6868	4b	1.00	20
	NCTC 7973	1a	0.19	53
	NCTC 5412	4b	0.90	13
	F4642	4b	0.92	14
	NCTC10357	1a	0.92	20
	BL87/41	4	0.66	25
	NCTC 5348	2	0.10	78
	SLCC2373	3a	1.20	17
	SLCC2540	3b	0.19	60
	SLCC2479	3c	0.15	60
	SLCC2374	4a	0.54	30
	SLCC2376	4c	0.19	90
	SLCC2377	4d	0.08	90
	SLCC2378	4e	0.56	28
	SLCC2482	7	0.45	36
	L3056	1/2a	0.49	30
	L4203	1/2a	0.36	41
	L4490	1/2b	0.29	55
	L1378	1/2b	0.09	150
	L4281	1/2c	0.11	120
	L3304	1/2c	0.12	90
<i>Listeria innocua</i>	L3253	4bx	0.66	26
	L2248	4bx	0.08	72
<i>Listeria ivanovii</i>	NCTC11288	6a	0.90	12
	NCTC11289	6a	0.69	22
<i>Listeria seeligeri</i>	NCTC11007		0.95	18
	SLCC5579		0.51	30
<i>Listeria velshimeri</i>	NCTC11856		1.10	15
	NCTC11857		0.29	36

TABLE 1: (continued)

ACTIVITY OF CLONED LYSIN AGAINST LISTERIA SPECIES				
Organism	Strain	Serotype	Relative <sup>a</sup> Activity	Time (min) <sup>b</sup> $\Delta OD_{600}=1$
<i>Listeria murrayi</i>	MCTC10812		0.86	15
<i>Listeria grayi</i>	NCTC10815		0.93	12
<i>Kurthia zopfii</i>	NCTC10597		0.54	28

[0040] Table 1 shows the relative sensitivity of a selection of strains of *Listeria* and *Kurthia zopfii* to the bacteriophage  $\phi$ LM4 lysin. a) Relative activity is the fall in optical density ( $OD_{600}$ ) from 2.3 achieved in 30 minutes divided by the equivalent fall obtained using *Listeria monocytogenes* F6868. b) The time (min) taken for a fall in optical density of  $OD_{600}$  from 2.3 to 1.3 ( $OD_{600}$  fall of 1) is recorded. Other strains tested which show no sensitivity to lysin were *Aeromonas hydrophila* NCTC 8049, *Bacillus cereus* NCTC 11143, *Brocothrix thermosphacta* NCTC 10822, *Carnobacterium piscicola* BL90/14, *Enterococcus faecalis* BL90/11, *Escherichia coli* BL90/12, *Klebsiella pneumoniae* NCFB 711, *Pseudomonas fluorescens* BL 78/45, *Staphylococcus aureus* NCTC 10652, *Streptococcus pneumoniae* NCTC 7465, *Streptococcus pyogenes* NCTC 2381.

[0041] In addition it was observed that the lysin was active at temperatures as low as 2°C. At 2°C addition of lysin to suspensions of *Listeria monocytogenes* caused a decrease of between 0.7 and 2.0  $OD_{600}$  units within 24 hours.

## EXAMPLE 2: USE OF LYSIN TO CONTROL LISTERIA

### Use as a free lysin

[0042] There are two distinct application concepts. One exploits a preparation of lysin enzyme manufactured by fermentation of a genetically engineered micro-organism that expresses the lysin gene product (Free lysin). The host organism may be *E. coli*, or any other bacterial species such as *Lactococcus lactis*, a yeast such as *Saccharomyces cerevisiae* or *Kluyveromyces fragilis* or a filamentous fungus such as *Aspergillus niger*. The lysin gene may be expressed intracellularly in which case a preparation may consist of a cell free lysate of the producing organism with some purification of the lysin, for example by ammonium sulphate precipitation and/or column chromatography. Alternatively the fermentation micro-organism may secrete lysin into the culture medium in which case the supernatant of the centrifuged fermentation broth provides the basis of a preparation, which again may require some purification.

[0043] The effectiveness of a crude extract of cloned *Listeria* lysin was demonstrated by its addition to skimmed milk containing *Listeria monocytogenes*. As illustrated in Fig. 9 the lysin preparation reduces the viable count of *Listeria monocytogenes* and after 22 days incubation at 8°C there is a viable count difference of  $10^8$  *Listeria* cfu between milk containing lysin and the control sample.

### Expression of Lysin by a genetically engineered micro-organism

[0044] An alternative application concept is to use a genetically engineered micro-organism that is compatible with a food or agricultural environment such as a species of lactic acid bacteria. Such an organism then grows in a food or agricultural environment and expresses an introduced gene for *Listeria* bacteriophage lysin. The gene may be expressed intracellularly and released into food or an agricultural environment by autolysis or induced lysis of that micro-organism. Alternatively the lysin may be secreted by a micro-organism so that active lysin is released into a food or agricultural environment by that viable micro-organism. In these cases the lysin gene is placed downstream of an appropriate promoter such as the lactose operon promoter or the proteinase promoter of *Lactococcus lactis* NCFB 712. Secretion may be achieved by fusion of the lysin structural gene to a known N terminal secretory leader such as those of the proteinase gene, the *usp45* gene or the nisin precursor gene of *Lactococcus lactis*. Suitable organisms for this application concept include strains of *Lactococcus lactis* in cheese and dairy products and *Lactobacillus plantarum* or *Pediococcus* species in agricultural silage.

[0045] The *Listeria* lysin gene from plasmid pF1328 was isolated together with its own ribosome binding site using the polymerase chain reaction. This fragment was cloned into the PstI site of *E. coli* vector pUC19 in both orientations (plasmids pF1531 and pF1532). Expression of this gene in *E. coli* strains was observed from one orientation only, under the control of the *lac*  $\alpha$  promoter of the vector (plasmid pF1531). Enzyme activity of cell extracts of this strain was comparable to that of *E. coli* strains carrying plasmid pF1322. Using plasmid pF1532 that did not express the lysin gene and cloning the lactococcal *lacA* promoter/*lacR* gene on a *Bam*HI fragment (Van Rooijen et al, (1992) *J. Bacteriol.* 174: 2273-2280) upstream of the lysin gene (plasmid pF1533) expression in *E. coli* of  $\phi$ LM-4 lysin from the lactococcal *lacA* promoter was obtained. The lytic activity of extracts from these *E. coli* strains was lower when the lysin gene was



expressed from the lacA promoter. The *SstI/SphI* fragment of pF1533 containing the  $\phi$ LM-4 lysin gene with the lacA promoter/lacR gene was cloned into the *SstI/SphI* sites of the lactococcal vector pTG262 (Shearman *et al* (1989) *Molecular and General Genetics* 218: 214-221) and the resulting plasmid pF1534 was used to transform *L. lactis* MG5267. As shown in Figure 10 cell extracts of this strain expressed  $\phi$ LM-4 lysin activity when grown on lactose; on glucose enzyme activity of cell extracts was reduced.

[0046] The  $\phi$ LM-4 lysin gene together with the lacA promoter/lacR gene was cloned into pF145, a plasmid expressing the *Lactococcus* phage  $\phi$ vML3 lysin gene which causes lysis during stationary phase of *L. lactis* cultures carrying the plasmid (Shearman *et al* (1992) *Biotechnology* 10: 196-199). The resulting plasmid pF1535 in *L. lactis* MG5267 when grown on lactose produced a culture that grew to stationary phase, then lysed as a consequence of the  $\phi$ vML3 lysin, releasing  $\phi$ LM-4 lysin into the culture supernatant.

### EXAMPLE 3: SPECIFIC DETECTION OF MICRO-ORGANISMS

[0047] The specificity of a bacteriophage lysin provides an opportunity to specifically detect those micro-organisms which are susceptible to it. For example to detect *Listeria* sp. the lysin described here may conveniently be used at a post enrichment stage where a broth culture of those micro-organisms present in a test sample is first produced. The identity of species of bacteria in the sample at this stage is unknown. The bacterial culture may be centrifuged and resuspended in an assay buffer (eg the one used here in studies of lysin specificity). A control preparation and separately a preparation containing active *Listeria* lysin are then added. Sufficient units of lysin activity are used to provide very effective lysis of any lysin susceptible cells (ie *Listeria*). After incubation for a short period (eg 30 min) any *Listeria* present will lyse, but other species will not. The presence of *Listeria* will then be detected by the lysis of bacteria in the sample treated with the lysin whereas no lysis occurs in the control.

[0048] The detection of lysis may be achieved by assaying an intracellular enzyme or metabolite. Especially useful enzyme assays are for phosphatase or for esterase. Alkaline phosphate can be assayed spectrophotometrically by following appearance of p-nitrophenol, which is yellow, from the colourless substrate p-nitrophenyl-phosphate at 405nm. Esterase activity can be assayed using fluorescein diacetate which is cleaved to acetate and fluorescent fluorescein and measuring the latter in a fluorometer. One especially suitable metabolite assay involves ATP detection. For this the well established luciferase assay in which ATP molecules generate light is exploited. Light emission may be measured in a luminometer. (An example of an end point detection reagent using luciferase-luciferin is marketed by Sigma Chemical Company as product L-1761).

### EXAMPLE 4: CLOSTRIDIUM TYROBUTYRICUM BACTERIOPHAGE $\phi$ P1 LYSIN

[0049] Bacteriophage  $\phi$ P1 was isolated from a landfill core sample using *Clostridium tyrobutyricum* NCFB 1755 as host. Bacteriophage  $\phi$ P1 was tested against six more strains of *C. tyrobutyricum*. Strains NCFB 1753 and NCFB 1756 supported the growth of bacteriophage and they were thus host strains as was the strain NCFB 1755. Against *C. tyrobutyricum* strains NCFB 1715, NCFB 1754, NCFB 1757 and NCFB 1790 an undiluted  $\phi$ P1 stock suspension gave a clear zone but diluting out did not result in individual bacteriophage plaques. This indicates that these strains were lysin sensitive but not bacteriophage sensitive. Bacteriophage  $\phi$ P1 thus produces a lysin with a broad specificity for strains of *C. tyrobutyricum*. Similar tests of bacteriophage  $\phi$ P1 with a wide variety of other bacteria showed no effect of the lysin or bacteriophage particles against *C. sporogenes* strains ATCC 17886, NCFB 1789, NCFB 1791; *C. butyricum* strains NCFB 1713, NCFB 857; *Lactobacillus buitteri* strains NCFB 110, F3327; *L. brevis* strains NCFB 1749, F3328; *L. helveticus* strains NCFB 1243, CNRZ 832; *L. bulgaricus* CNRZ448; *L. plantarum* strains NCFB 1752, NCFB 82, NCFB 963; *Escherichia coli* BL 90/12; *Bacillus cereus* NCTC 1143.

[0050] Bacteriophage  $\phi$ P1 was deposited at the National Collections of Industrial and Marine Bacteria, 23 St Machar Drive, Aberdeen, AB2 1RY, Scotland on 5 April 1991 and a new deposit was made on 4 July 1991 under the Budapest Treaty and has been accorded Accession No NCIMB 40400.

### Claims

1. A formulation comprising a lysin of a bacteriophage of *Clostridia* spp., or a variant thereof having an amino acid sequence which is substantially identical to such a lysin, substantially free of the bacteriophage itself.
2. A substantially pure preparation of a lysin of a bacteriophage of *Clostridia* spp.
3. A formulation according to Claim 1 or a preparation according to Claim 2 wherein the bacteriophage is a bacteriophage of *Clostridia tyrobutyricum*.

## EP 1 300 082 A2

4. A formulation according to Claim 1 or a preparation according to Claim 2 wherein the bacteriophage is øP1.
5. A formulation according to Claim 1 or a preparation according to Claim 2 wherein the bacteriophage is øP1, deposited under the Budapest Treaty under accession No NCIMB 40400.
6. A nucleotide coding sequence for a lysin as defined in any one of Claims 2 to 5.
7. An expression vehicle comprising a coding sequence according to Claim 6 and regulatory regions associated therewith for expression of the coding sequence in a suitable host.
8. A microbial host transformed with means comprising the coding sequence of Claim 6 to express a lysin of a bacteriophage of *Clostridia* spp.
9. A host according to Claim 8 which is a food grade microorganism.
10. A lysin derived from the cultivation of a host according to Claim 8 or 9.
11. A lysin as defined in any one of Claims 1 to 5 or Claim 10 for use as an antimicrobial agent.
12. Use of a lysin as defined in any one of Claims 1 to 5 or Claim 10 to destroy a food-contaminating bacterium prejudicial to health in or on food products.
13. A method of destroying micro-organisms characterised in that said micro-organisms are lysed with a formulation according to any one of Claims 1 to 5 or Claim 10.
14. A method according to Claim 13 wherein pathogenic strains of *Clostridia* are destroyed in or on food products.
15. A method of testing for *Clostridia* in a sample, comprising adding a bacteriophage lysin substantially free of intact bacteriophage to the sample and determining whether bacterial cells have been lysed thereby, wherein the lysin is a lysin of Clostridial bacteriophage or a variant thereof having an amino acid sequence which is substantially identical to such a lysin.
16. A method according to Claim 15 wherein the bacteriophage is a *Clostridium tyrobutyricum* bacteriophage
17. A method according to Claim 16 wherein the bacteriophage is øP1.




Fig. 1




FIGURE 2




FIGURE 3




Fig. 4




FIGURE 5

```

1  AAGCTTTACT GGATAACGAC AAACCTAGAAC TGACAGATGA AGGACTGAAA
51  GGCCTTGACG AACAGCTAGG AGCATTGCAA GAAAGCGATG CTTATTTGTT
101 TGCTCAAGAA AGCGAGGCGG GCGGGAAGTG GTTTATTTGA ACAATTTACT
151 AATCGAGCTA AAAAATATGG AAAGGATGAT TAATAATGGC ATTAACAGAG
201 GCATGGCTAA TTGAAAAAGC AAATCGCAAA TTGAATACGT CAGGTATGAA
251 TAAAGCTACA TCTGATAAGA CTCGGAATGT AATTAAAAAA ATGGCAAAAG
301 AAGGGATTTA TCTTTGTGTT GCGCAAGGTT ACCGCTCAAC AGCGGAACAA
351 AATGCGCTAT ATGCACAAGG GAGAACCAAA CCTGGAGCGA TTGTTACTAA
401 TGCTAAAGGT GGGCAATCTA ATCATAATTT CGGTGTAGCA GTTGATTTGT
451 GCTTGTATAC GAGCGACGGA AAAGATGTTA TTTGGGAGTC GACAACTTCC
501 CGGTGGAAAA AGGTTGTTGC TGCTATGAAA GCGGAAGGAT TCGAATGGGG
551 CGGAGATTGG AAAAGTTTTA AAGACTATCC GCATTTTGAA CTATGTGACG
601 CTGTAAGTGG TGAGAAAATC CCTACTGCGA CACAAAACAC CAATCCAAAC
651 AGACATGATG GGAAAATCGT TGACAGCGCG CCACTATTGC CAAAAATGGA
701 CTTTAAATCA AATCCAGCGC GCATGTATAA ATCAGGAACT GAGTTCCTAG
751 TATATGAACA TAATCAATAT TGGTACAAGA CGTACATCAA CGACAAATTA
801 TACTACATGT ATAAGAGCTT TTGCGATGTT GTAGCTAAAA AAGATGCAAA
851 AGGACGCATC AAAGTTCGAA TTAAAAGCGC GAAAGACTTA CGAATTCCAG
901 TTTGGAATAA CACAAAATTG AATTCTGGGA AAATTAAATG GTATGCACCC
951 AATACAAAAT TAGCATGGTA CAACAACGGA AAAGGATACT TGGAACCTG
1001 GTATGAAAAG GATGGCTGGT ACTACACAGC GAACTACTTC TTAAAATAAA
1051 AAGTCCCGGT TTGAGCTGGG CTTTTTATTT TGAAAGTGAC TAACAAAAAA
1101 TGTAATAAAA ATGTAATAAT CCAAGTAAGT TGTATAAAAT TTGCAGAATT
1151 AGAACGTTTT ATTGATAAAT ACACCTTATG AAGAGTATTT TTGGCTATAT
1201 TTGCGCATTa TAGGGTTGAA TGTAACACTA TATGTAGAAT TC

```

FIGURE 6






FIGURE 7

MetAlaLeuThrGluAlaTrpLeuIleGluLysAlaAsnArgLysLeuAsnThrSerGly  
 ATGGCATTAAACAGAGGCATGGCTAATTGAAAAAGCAAATCGCAAATTGAATACGT CAGGT 245  
 186 -----+-----+-----+-----+-----+-----+-----  
 TACCGTAATTGTCTCCGTACCGATTAAC TTTTCGTTTAGCGTTAACTTATGCAGTCCA  
  
 MetAsnLysAlaThrSerAspLysThrArgAsnValIleLysLysMetAlaLysGluGly  
 ATGAATAAAGCTACATCTGATAAGACTCGGAATGTAATTAATAAATAATGGCAAAAGAAGGG 305  
 246 -----+-----+-----+-----+-----+-----+-----  
 TACTTATTTTCGATGTAGACTATTCTGAGCCTTACATTAATTTTTTTTACCGTTTTCTTCCC  
  
 IleTyrLeuCysValAlaGlnGlyTyrArgSerThrAlaGluGlnAsnAlaLeuTyrAla  
 ATTTATCTTTGTGTTGCGCAAGGTTACCGCTCAACAGCGGAACAAAATGCGCTATATGCA 365  
 306 -----+-----+-----+-----+-----+-----+-----  
 TAAATAGAAACACAACGCGTTCCAATGGCGAGTTGTCGCCTTGTTTTACGCGATATACGT  
  
 GlnGlyArgThrLysProGlyAlaIleValThrAsnAlaLysGlyGlyGlnSerAsnHis  
 CAAGGGAGAACCAAACCTGGAGCGATTGTTACTAATGCTAAAGGTGGGCAATCTAATCAT 425  
 366 -----+-----+-----+-----+-----+-----+-----  
 GTTCCCTCTTGGTTTGGACCTCGCTAACAAATGATTACGATTTCCACCGTTAGATTAGTA  
  
 AsnPheGlyValAlaValAspLeuCysLeuTyrThrSerAspGlyLysAspValIleTrp  
 AATTCGGTGTAGCAGTTGATTTGTGCTTGATACGAGCGACGAAAAGATGTTATTTGG 485  
 426 -----+-----+-----+-----+-----+-----+-----  
 TTAAAGCCACATCGTCAACTAAACACGAACATATGCTCGCTGCCTTTTCTACAATAAACC  
  
 GluSerThrThrSerArgTrpLysLysValValAlaAlaMetLysAlaGluGlyPheGlu  
 GAGTCGACAACCTCCCGGTGGAAAAGGTTGTTGCTGCTATGAAAGCGGAAGGATTCGAA 545  
 486 -----+-----+-----+-----+-----+-----+-----  
 CTCAGCTGTTGAAGGGCCACCTTTTTCCAACAACGACGATACTTTTCGCCTTCCTAAGCTT  
  
 TrpGlyGlyAspTrpLysSerPheLysAspTyrProHisPheGluLeuCysAspAlaVal  
 TGGGGCGGAGATTGGAAAAGTTTTAAAGACTATCCGCATTTTGAAGTATGTGACGCTGTA 605  
 546 -----+-----+-----+-----+-----+-----+-----  
 ACCCGCCTCTAACCTTTTCAAATTTCTGATAGGCGTAAAACCTTGATACACTGCGACAT  
  
 SerGlyGluLysIleProThrAlaThrGlnAsnThrAsnProAsnArgHisAspGlyLys  
 AGTGGTGAGAAAATCCCTACTGCGACACAAAACACCAATCCAAACAGACATGATGGGAAA 665  
 606 -----+-----+-----+-----+-----+-----+-----  
 TCACCACTCTTTTAGGGATGACGCTGTGTTTGTGGTTAGGTTTGTCTGTACTACCCTTT  
  
 IleValAspSerAlaProLeuLeuProLysMetAspPheLysSerAsnProAlaArgMet  
 ATCGTTGACAGCGCGCCACTATTGCCAAAATGGACTTTAAATCAAATCCAGCGCGCATG 725  
 666 -----+-----+-----+-----+-----+-----+-----  
 TAGCAACTGTGCGCGGTGATAACGGTTTTTACCTGAAATTTAGTTTAGGTGCGCGGTAC  
  
 TyrLysSerGlyThrGluPheLeuValTyrGluHisAsnGlnTyrTrpTyrLysThrTyr  
 TATAAATCAGGAAGTGAAGTTCTTAGTATATGAACATAATCAATATTGGTACAAGCGTAC 785  
 726 -----+-----+-----+-----+-----+-----+-----  
 ATATTTAGTCCTTGACTCAAGAATCATATACTTGTATTAGTTATAACCATGTTCTGCATG

FIGURE 8 (START)

	IleAsnAspLysLeuTyrTyrMetTyrLysSerPheCysAspValValAlaLysLysAsp	
	ATCAACGACAAATTATACTACATGTATAAGAGCTTTTGCGATGTTGTAGCTAAAAAGAT	
786	-----+-----+-----+-----+-----+-----+-----	845
	TAGTTGCTGTTTAATATGATGTACATATTCTCGAAAACGCTACAACATCGATTTTTCTA	
	AlaLysGlyArgIleLysValArgIleLysSerAlaLysAspLeuArgIleProValTrp	
	GCAAAAGGACGCATCAAAGTTCAATTAAAAGCGCGAAAGACTTACGAATTCAGTTTGG	
846	-----+-----+-----+-----+-----+-----+-----	905
	CGTTTTCTGCGTAGTTTCAAGCTTAATTTTCGCGCTTTCTGAATGCTTAAGGTCAAACC	
	AsnAsnThrLysLeuAsnSerGlyLysIleLysTrpTyrAlaProAsnThrLysLeuAla	
	AATAACACAAAATTGAATTCTGGGAAAATTAAATGGTATGCACCCAATACAAAATTAGCA	
906	-----+-----+-----+-----+-----+-----+-----	965
	TTATTGTGTTTTAACTTAAGACCCTTTTAATTTACCATACGTGGGTTATGTTTTAATCGT	
	TrpTyrAsnAsnGlyLysGlyTyrLeuGluLeuTrpTyrGluLysAspGlyTrpTyrTyr	
	TGGTACAACAACGGAAAAGGATACTTGGAACCTCTGGTATGAAAAGGATGGCTGGTACTAC	
966	-----+-----+-----+-----+-----+-----+-----	1025
	ACCATGTTGTTGCCTTTTCCTATGAACCTTGAGACCATACTTTTCCTACCGACCATGATG	
	ThrAlaAsnTyrPheLeuLys	
	ACAGCGAACTACTTCTTAAAA	
1026	-----+-----+-----+-----+-----+-----+-----	1046
	TGTCGCTTGATGAAGAATTTT	

FIGURE 8 (END)




FIGURE 9




FIGURE 10



(12) **EUROPEAN PATENT APPLICATION**

(88) Date of publication A3:  
**19.11.2003 Bulletin 2003/47**

(43) Date of publication A2:  
**09.04.2003 Bulletin 2003/15**

(21) Application number: **02079804.7**

(22) Date of filing: **21.04.1992**

(51) Int Cl.<sup>7</sup>: **A23K 1/17, A23L 3/3463,**  
**A23L 3/3571, C12N 1/06,**  
**C12N 1/15, C12N 1/19,**  
**C12N 1/20, C12N 7/01,**  
**C12N 9/14, C12N 15/33,**  
**C12N 15/55, C12N 15/74,**  
**C12Q 1/04, C12Q 1/10,**  
**C12Q 1/18, C12Q 1/34,**  
**C12Q 1/70**  
**// C12R1:145**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GR IT LI LU MC NL PT SE**

(30) Priority: **20.04.1991 GB 9108498**

(62) Document number(s) of the earlier application(s) in  
accordance with Art. 76 EPC:  
**92303533.1 / 0 510 907**

(71) Applicant: **THE INSTITUTE OF FOOD RESEARCH**  
**Norwich NR4 7UA (GB)**

(72) Inventor: **Gasson; Michael John**  
**Dereham, Norfolk NR20 5NA (GB)**

(74) Representative: **Thomas, Philip John Duval et al**  
**Eric Potter Clarkson,**  
**Park View House,**  
**58 The Ropewalk**  
**Nottingham NG1 5DD (GB)**

(54) **Lysin of a bacteriophage of Clostridia spp**

(57) Bacteriophages of food-contaminating or pathogenic bacteria or the lysins thereof are used to kill such bacteria. Examples include lysins from bacteriophages of *Listeria monocytogenes* and *Clostridium tyrobutyricum*.

Tests for bacterial contamination can be made specific for specific bacteria by using the appropriate bacteriophage or lysin thereof and determining whether cells are lysed thereby.




Fig. 1



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 02 07 9804

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1982 EFIMOVA M G: "ISOLATION OF PURIFIED TYPE A CLOSTRIDIUM-PERFRINGENS THETA HEMO LYSIN AND ITS USE FOR THE TITRATION OF TOXINS AND SERA MICE TOXICITY EQUINE ANTI SERUM ANTI TOXIN BINDING ANTI TOXIN QUANTITATION" Database accession no. PREV198325053312 XP002239110 * abstract *	1-5,10, 11	A23K1/17 A23L3/3463 A23L3/3571 C12N1/06 C12N1/15 C12N1/19 C12N1/20 C12N7/01 C12N9/14 C12N15/33 //C12R1:145
Y	& ZHURNAL MIKROBIOLOGII EPIDEMIOLOGII I IMMUNOBIOLOGII, no. 10, 1982, ISSN: 0372-9311 ---	6-9	
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1982 YOSHINO S ET AL: "SOME PROPERTIES OF AUTO LYSIN OF CLOSTRIDIUM-SACCHAROPERBUTYLACETONICUM" Database accession no. PREV198375036608 XP002239111 * abstract *	1-5,10, 11	TECHNICAL FIELDS SEARCHED (Int.Cl.7)  A23L C12N C12Q
Y	& AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 46, no. 5, 1982, pages 1243-1248, ISSN: 0002-1369 --- -/--	6-9	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 11 September 2003	Examiner Thiele, U
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons --- &: member of the same patent family, corresponding document	

EPO FORM 1503 03.02 (P4/C01)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 02 07 9804

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL1.7)
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1982 YOSHINO S ET AL: "PECULIAR BEHAVIOR OF AUTO LYSIN OF CLOSTRIDIUM-SACCHAROPERBUTYLACETO NICUM ON ANION EXCHANGERS" Database accession no. PREV198375013418 XP002239112 * abstract *	1-5,10, 11	
Y	& SCIENCE BULLETIN OF THE FACULTY OF AGRICULTURE KYUSHU UNIVERSITY, vol. 36, no. 2-3, 1982, pages 115-122, ISSN: 0368-6264 ---	6-9	
X	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1981 DOLGIKH M S ET AL: "ISOLATION OF THETA HEMO LYSIN FROM CLOSTRIDIUM-PERFRINGENS" Database accession no. PREV198273076511 XP002239113 * abstract *	1-5,10, 11	
Y	& VOPROSY MEDITSINSKOI KHIMII, vol. 27, no. 1, 1981, pages 129-132, ISSN: 0042-8809 ---	6-9	
X	HIRANO S ET AL: "A PHAGE-ASSOCIATED LYSIN OF CLOSTRIDIUM PERFRINGENS" ACTA MEDICA UNIVERSITATIS KAGOSHIMAENSIS, KAGOSHIMA, JP, vol. 14, no. 1, 1972, pages 91-101, XP000650442 ISSN: 0001-611X ---	1-5,10, 11	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
Y	* the whole document * ---	6-9	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 11 September 2003	Examiner Thiele, U
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document	

EPO FORM 1503 (03.02) (P4/C01)





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 02 07 9804

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Y	BOIZET B ET AL: "CLONING, EXPRESSION AND SEQUENCE ANALYSIS OF AN ENDOLYSIN-ENCODING GENE OF LACTOBACILLUS BULGARICUS BACTERIOPHAGE MV1" GENE, ELSEVIER BIOMEDICAL PRESS. AMSTERDAM, NL, vol. 94, 1990, pages 61-67, XP001019271 ISSN: 0378-1119 * the whole document *	6-9	
Y	GARCIA J L ET AL: "CLONING, PURIFICATION, AND BIOCHEMICAL CHARACTERIZATION OF THE PNEUMOCOCCAL BACTERIOPHAGE CP-1 LYSIN" JOURNAL OF VIROLOGY, NEW YORK, US, US, vol. 61, no. 8, 1 August 1987 (1987-08-01), pages 2573-2580, XP000108167 ISSN: 0022-538X * the whole document *	6-9	
A	DATABASE MEDLINE 'Online! US NATIONAL LIBRARY OF MEDICINE (NLM), BETHESDA, MD, US; April 1991 (1991-04) BOGOVAZOVA G G ET AL: "The efficacy of Klebsiella pneumoniae bacteriophage in the therapy of experimental Klebsiella infection!" Database accession no. NLM1882608 XP002239114 * abstract * & ZHURNAL MIKROBIOLOGII, EPIDEMIOLOGII, I IMMUNOBIOLOGII. USSR APR 1991, no. 4, April 1991 (1991-04), pages 5-8, ISSN: 0372-9311 ----- -/--		
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 11 September 2003	Examiner Thiele, U
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P4/C01)



European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 02 07 9804

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
T	<p>JOERGER R D: "Alternatives to antibiotics: Bacteriocins, antimicrobial peptides and bacteriophages." POULTRY SCIENCE, 'Online! vol. 81, no. Supplement 1, 2002, page 32 XP002239109 91st Annual Meeting of the Poultry Science Association; Newark, DE, USA; August 08-11, 2002 ISSN: 0032-5791 Retrieved from the Internet: &lt;URL:ww.poultryscience.org/meet/91st/psabs12.pdf&gt; 'retrieved on 2003-04-23! * the whole document *</p> <p>---</p>		
X	<p>NAKAMURA S ET AL: "CLOSTRIDIUM-PERFRINGENS SPECIFIC LYSIN" CANADIAN JOURNAL OF MICROBIOLOGY, vol. 23, no. 5, 1977, pages 601-606, XP002246438 ISSN: 0008-4166 *sections headed: "Introduction", "Discussion"; page 605, lines 7 - 10*</p> <p>---</p>	15-17	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	<p>NAKAMURA S ET AL: "LECITHINASE NEGATIVE VARIANTS OF CLOSTRIDIUM-PERFRINGENS THE IDENTITY OF CLOSTRIDIUM-PLAGARUM WITH CLOSTRIDIUM-PERFRINGENS" CANADIAN JOURNAL OF MICROBIOLOGY, vol. 22, no. 10, 1976, pages 1497-1501, XP009017157 ISSN: 0008-4166 * abstract *</p> <p>-----</p>	15-17	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 11 September 2003	Examiner Thiele, U
CATEGORY OF CITED DOCUMENTS		<p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>-----</p> <p>&amp; : member of the same patent family, corresponding document</p>	
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p>			

EPO FORM 1503 03.02 (PSC01)



European Patent  
Office

Application Number  
EP 02 07 9804

#### CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

#### LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

- ☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent  
Office

LACK OF UNITY OF INVENTION  
SHEET B

Application Number  
EP 02 07 9804

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims: 1-14

Formulation comprising a lysin of a bacteriophage of *Clostridia* spp., in particular *Clostridia tyrobutyricum* and phiP1, or a variant thereof, substantially pure preparation of said lysin, destruction of microorganisms using these lysins, destruction of bacteria, in particular *Clostridium*, on food using these lysins, DNA encoding these lysins, expression vehicles and hosts to express these lysins, recombinant lysin

2. Claims: 15-17

Method for testing for *Clostridia* in a sample by adding a lysin of *Clostridia* bacteriophage, in particular *Clostridium tyrobutyricum* and phiP1, or a variant thereof, to the sample